

May 10, 2003

IMPROVEMENT OF THE THEOREM ON LOCAL ERGODICITY

Nándor Simányi¹

University of Alabama at Birmingham
 Department of Mathematics
 Campbell Hall, Birmingham
 AL 35294 U.S.A.
 E-mail: simanyi@math.uab.edu

Abstract. We prove here that in the Theorem on Local Ergodicity for Semi-Dispersive Billiards (proved by N. I. Chernov and Ya. G. Sinai in 1987) the recently added condition (by P. Bálint, N. Chernov, D. Szász, and I. P. Tóth, in order to save this fundamental result) on the algebraic character of the smooth boundary components of the configuration space is unnecessary. Having saved the theorem in its original form by using additional ideas in the spirit of the initial proof, the result becomes stronger and it applies to a larger family of models.

Primary subject classification: 37D50

Secondary subject classification: 34D05

§1. INTRODUCTION

Semi-Dispersive Billiards. Assume that $\mathbf{Q} \subset \mathbb{R}^d$ or $\mathbf{Q} \subset \mathbb{T}^d = \mathbb{R}^d/\mathbb{Z}^d$ ($d \geq 2$) is an open and connected subset with a non-empty, piecewise smooth (i. e. piecewise C^3) boundary $\partial\mathbf{Q}$ and compact closure $\overline{\mathbf{Q}} = \mathbf{Q} \cup \partial\mathbf{Q}$. Throughout this paper the word “smooth” means class C^3 for submanifolds of $\mathbf{M}$, whereas its meaning is C^2 for submanifolds of $\partial\mathbf{M}$. In order to avoid unnecessary technical complications, we

¹Research supported by the National Science Foundation, grant DMS-0098773.

also assume that the d -dimensional spatial angle $\angle(q)$ subtended by $\mathbf{Q}$ at any of its boundary points $q \in \partial\mathbf{Q}$ is positive. We also assume that the smooth components $\partial\mathbf{Q}_i$ ($i = 1, 2, \dots, k$) of $\partial\mathbf{Q}$ are (not necessarily strictly) concave, i. e. they are “bending away” from $\mathbf{Q}$. In technical terms this means the following: If we supply the smooth component $\partial\mathbf{Q}_i$ with the field $\nu(q)$ ($q \in \partial\mathbf{Q}_i$) of unit normal vectors pointing inward $\mathbf{Q}$, then the second fundamental form $K(q)$ of $\partial\mathbf{Q}_i$ at q (with respect to the selected field of unit normal vectors) is a non-negative operator for any $q \in \partial\mathbf{Q}_i$ and $i = 1, \dots, k$. Finally, one always assumes that at any boundary point $q \in \partial\mathbf{Q}_i \cap \partial\mathbf{Q}_j$ ($1 \leq i < j \leq k$) the tangent vectors of $\mathbb{R}^d$ (or $\mathbb{T}^d$) at q pointing inward $\mathbf{Q}$ form a convex cone.

The semi-dispersive billiard system (flow) $(\mathbf{M}, \{S^t\}_{t \in \mathbb{R}}, \mu)$ is a dynamical system describing the uniform motion (at a unit speed) of a point particle inside the flat domain $\mathbf{Q}$ that bounces back at $\partial\mathbf{Q}$ in a fully elastic manner, according to the law of geometric optics, that is, the angle of reflection equals the angle of incidence. The phase space $\mathbf{M}$ of the arising flow is essentially the unit tangent bundle $\overline{\mathbf{Q}} \times \mathbb{S}^{d-1}$ ($\mathbb{S}^{d-1}$ being the unit sphere of velocities) modulo the natural identification of the incoming (pre-collision) and outgoing (post-collision) velocities at the boundary $\partial\mathbf{M} = \partial\mathbf{Q} \times \mathbb{S}^{d-1}$. The flow $\{S^t\}_{t \in \mathbb{R}}$ is the time-evolution of the system, and the (invariant) Liouville measure μ turns out to be the normalized Lebesgue measure: $d\mu = \text{const} \cdot dq \cdot dv$, where $x = (q, v) \in \mathbf{M}$, $q \in \mathbf{Q}$, $v \in \mathbb{S}^{d-1}$ is the standard expansion of a phase point x into the configuration and velocity components.

In the case $K(q) > 0$ (for every $q \in \partial\mathbf{Q}$) we speak about a “dispersive billiard system”.

The primary examples of semi-dispersive billiards are the so called cylindric billiards (see [S-Sz(2000)]), among which the most important special models are the hard ball systems, see §2 of [S-Sz(1999)]. For a more detailed introduction to these dynamical systems, please see §2 (the sections containing the prerequisites) of the papers [K-S-Sz(1990)], [S-Sz(1999)], and [Sim(2002)].

Assume that $(\mathbf{M}, \{S^t\}_{t \in \mathbb{R}}, \mu)$ is a semi-dispersive billiard flow with the following additional properties:

(1) For ν -almost every singular phase point $x \in \mathcal{SR}^+ (\subset \partial\mathbf{M})$ the forward semi-trajectory $S^{[0, \infty)}x$ of x is sufficient (or *geometrically hyperbolic*, see §2 for $\mathcal{SR}^+$ and the notion of sufficiency). This condition is precisely Condition 3.1 of [K-S-Sz(1990)];

(2) (Regularity of the set of degenerate tangencies, a slightly strengthened version of Condition 3.2 of [K-S-Sz(1990)]) The set

$$DT = \{x = (q, v) \in \partial\mathbf{Q} \times \mathbb{S}^{d-1} \mid \langle v, n(q) \rangle = 0, \quad K(q)[v] = 0\}$$

of *degenerate tangencies* is a compact cell sub-complex (finite union of smooth, compact submanifolds) of $\partial\mathbf{M}$ with the property $\dim(DT) \leq 2d - 4 (= \dim(\partial\mathbf{M}) -$

2). This condition precisely means that the set $\{q \in \partial\mathbf{Q} \mid K(q) \equiv 0\}$ has an empty interior in $\partial\mathbf{Q}$, i. e. $\partial\mathbf{Q}$ has no flat piece.

(3) A given phase point $x_0 \in \mathbf{M} \setminus \partial\mathbf{M}$ has a sufficient orbit $S^{\mathbb{R}}x_0$ (see §2) with at most one (and simple) singularity on it.

Under the above conditions, the Theorem on Local Ergodicity claims that some open neighborhood U_0 of x_0 belongs (modulo the zero sets) to a single ergodic component of the flow $\{S^t\}_{t \in \mathbb{R}}$.

This fundamental result was originally proved by N. I. Chernov and Ya. G. Sinai as Theorem 5 in [S-Ch(1987)], being actually the corollary of the rather technical Lemma 3 there. Later on, the proof of Lemma 3 was somewhat clarified and put into a more general context as Theorem 3.6 of [K-S-Sz(1990)]. In that paper the exact counterpart of the above cited Theorem on Local Ergodicity is Corollary 3.12. Condition (1) is often called the “Chernov-Sinai Ansatz” in the literature.

Recently it was discovered in [B-Ch-Sz-T(2002)] that the original proof of the Theorem on Local Ergodicity contained a gap: The utmost important geometric inequality of Lemma 4.6 of [K-S-Sz(1990)] does not follow automatically from the geometry of the singularities of the billiard map. The paper [B-Ch-Sz-T(2002)] turned out to be a successful attempt — with beautiful mathematics — to fix the original proof at the expense of additionally assuming that the smooth components $\partial\mathbf{Q}_i$ of the boundary $\partial\mathbf{Q}$ are algebraic (i. e. polynomially defined) hypersurfaces, not only C^3 -smooth manifolds. However, all experts strongly believe that C^3 smoothness should be enough for the result. Also, due to this restriction, the scope of the Theorem on Local Ergodicity becomes somewhat narrower than expected, though the most important models (like hard ball systems, cylindric billiards) are still algebraic, and thus covered by the result of [B-Ch-Sz-T(2002)].

The main result of this paper is saving the Theorem on Local Ergodicity in its original form, without assuming the algebraic nature of the boundary components $\partial\mathbf{Q}_i$. This will be accomplished by obtaining a correct proof for the estimation of Lemma 4.6 of [K-S-Sz(1990)], see Lemma 3 below.

The value of the present paper is two-fold:

First, as we said above, the present additions to the original proof do not require the algebraicity and, this way, they give the theorem the expected range of applicability. The mentioned geometric inequality of Lemma 4.6 of [K-S-Sz(1990)] (proved here as Lemma 3) will follow from the new, more dynamical approach, despite the fact that — as it has been revealed in [B-Ch-Sz-T(2002)] — the second derivatives (the curvature tensor) of the singularity components K_i may blow up near the boundary ∂K_i (Here K_i is a smooth component of the singularity set $\mathcal{R}^{\leq n}$ of the n -th iterate of the billiard map), so that even the tangent hyperplane of K_i may be undefined at some points of ∂K_i . This wild geometry has the potential for

breaking down the geometric inequality of Lemma 4.6 of [K-S-Sz(1990)]. We will be presenting the new proof by introducing some interesting additions to the original ideas of Chernov and Sinai.

Secondly, the proof in [B-Ch-Sz-T(2002)] turns out to be a bit involved and technically difficult, though nice mathematics, in contrast with my new, simplified approach, which is based much more on dynamical ideas, rather than algebraic and geometric measure-theoretic ones. In this sense my approach lies closer to the spirit of the original Chernov-Sinai proof.

Since the improvement shown below is an addition to the original Chernov-Sinai proof, I did not find it appropriate to repeat the entire proof but, instead, I opted for presenting essentially the proof for Lemma 4.6 only, with frequent references to the papers [S-Ch(1987)] and [K-S-Sz(1990)].

§2. PREREQUISITES

Singularities and Trajectory Branches. The billiard system $(\mathbf{M}, \{S^t\}_{t \in \mathbb{R}}, \mu)$ has two types of singularities. The first one is caused by the presence of the so called “tangential reflections” $x_0 = (q_0, v_0) \in \partial\mathbf{M}$ for which $\langle v_0, n(q_0) \rangle = 0$, i. e. the velocity v_0 happens to lie inside the tangent hyperplane $\mathcal{T}_{q_0}\partial\mathbf{Q}$ of $\partial\mathbf{Q}$ at $q_0 \in \partial\mathbf{Q}$. If the trajectory $S^{(-\infty, \infty)}y$ of a phase point hits a tangential reflection at time $t_0 > 0$ ($S^{t_0}y = x_0$), then the billiard map $S^{t_0+\epsilon}$ is still continuous at y , though it ceases to be smooth. The often studied first return map $T : \partial\mathbf{M} \rightarrow \partial\mathbf{M}$ of the boundary $\partial\mathbf{M}$ is no longer even continuous at the inverse image $T^{-1}x_0$ of a phase point x_0 with tangential reflection.

The second type of singularity, the so called “multiple reflection” takes place when the flow $S^{t_0}y = x_0 = (q_0, v_0^-) \in \partial\mathbf{M}$ (v_0^- is the “incoming” velocity) hits more than one boundary components of $\mathbf{M}$, i. e. when $q_0 \in \partial\mathbf{Q}_i \cap \partial\mathbf{Q}_j$ with $1 \leq i < j \leq k$. We are going to briefly describe the discontinuity of the flow $\{S^t\}$ caused by a multiple collision at time t_0 . Assume first that the pre-collision velocity is given. What can we say about the possible post-collision velocity? Let us perturb the pre-collision phase point (at time $t_0 - 0$) infinitesimally, so that the collisions at $\sim t_0$ occur at infinitesimally different moments. By applying the collision laws to the arising finite sequence of collisions, (the finiteness follows from Theorem 1 of [B-F-K(1998)]) we see that on the two sides of the current (codimension-one) singularity manifold there are two, significantly different continuations (the so called branches) of the trajectory: On one side the finite sequence of collisions near time t_0 begins with a $\partial\mathbf{Q}_i$ -reflection, then it continues in an alternating manner with $\partial\mathbf{Q}_j$, $\partial\mathbf{Q}_i, \dots$ reflections all the way until the resulting (reflected) velocity v^+ happens to point inward $\mathbf{Q}$. On the other side of the singularity, however, the corresponding

finite, alternating sequence of reflections begins with a $\partial\mathbf{Q}_j$ -collision. These two continuations (branches) result in two (typically different) outgoing velocities v^+ . Analogous statements can be made about the backward continuations and the $v^+ \mapsto v^-$ correspondence. It follows from Lemma 4.1 of [K-S-Sz(1990)] that a typical singular phase point x_0 has the property that its trajectory hits a singularity only once, say, at time $t_0 > 0$ and

(a) $q(S^{t_0}x_0)$ lies only on one boundary component $\partial\mathbf{Q}_i$ if $S^{t_0}x_0$ is a tangential reflection;

(b) $q(S^{t_0}x_0)$ lies exactly on two boundary components $\partial\mathbf{Q}_i$ and $\partial\mathbf{Q}_j$ ($i \neq j$) if $S^{t_0}x_0$ is not a tangency;

(c) none of the arising two trajectory branches hits a singularity any more.

In this case we say that the trajectory of x_0 encounters a so called *simple singularity*. In this situation the entire orbit $S^{(-\infty, \infty)}x_0$ has exactly two branches.

Since, in the case of multiple collisions, there is no unique continuation of the trajectories, we need to make a clear distinction between the set of singular reflections $\mathcal{SR}^+$ supplied with the outgoing velocity v^+ , and the set of singular reflections $\mathcal{SR}^-$ supplied with the incoming velocity v^- . For typical phase points $x^+ \in \mathcal{SR}^+$ the forward trajectory $S^{[0, \infty)}x^+$ is non-singular and uniquely defined, and analogous statement holds true for typical phase points $x^- \in \mathcal{SR}^-$ and the backward trajectory $S^{(-\infty, 0]}x^-$. For a more detailed exposition of singularities and trajectory branches, the reader is kindly referred to §2 of [K-S-Sz(1990)]. We denote by $\mathcal{SR} = \mathcal{SR}^- \cup \mathcal{SR}^+$ the set of all singular reflections.

Sufficiency (Geometric Hyperbolicity). Let $S^{[a, b]}x_0$ be a non-singular (i. e. smooth) trajectory segment of a semi-dispersive billiard flow $(\mathbf{M}, \{S^t\}_{t \in \mathbb{R}}, \mu)$. At the starting phase point $y_0 = S^a x_0 = (q_0, v_0)$ we construct the submanifold

$$\Sigma_0 = \{(q, v_0) \mid q - q_0 \perp v_0 \text{ and } \|q - q_0\| < \epsilon_0\}$$

of $\mathbf{M}$ with a fixed number $\epsilon_0 \ll 1$. Denote by Σ_1 the connected component of the image $S^{b-a}(\Sigma_0)$ containing $S^b x_0 = S^{b-a} y_0$. It is easy to see that the smooth submanifold Σ_1 of $\mathbf{M}$ has the form

$$\Sigma_1 = \{(q, v(q)) \mid q \in \tilde{\Sigma}_1\},$$

where $\tilde{\Sigma}_1 \subset \mathbf{Q}$ is a smooth hypersurface of the configuration space $\mathbf{Q}$ supplied with the field $v(q)$ of unit normal vectors. A direct consequence of the semi-dispersive property is that the second fundamental form $K(q)$ of $\tilde{\Sigma}_1$ at any $q \in \tilde{\Sigma}_1$ (with respect to the above field of unit normal vectors) is non-negative. The trajectory segment $S^{[a, b]}x_0$ is said to be *sufficient* (or *geometrically hyperbolic*) if $K(q_1) > 0$, where $S^b x_0 = (q_1, v_1)$.

Definition. The trajectory segment $S^{[a,b]}x$ containing a simple singularity is said to be sufficient if and only if both branches of this trajectory segment are sufficient.

Definition. The phase point $x \in \mathbf{M}$ with at most one (and simple) singularity is said to be sufficient if and only if its whole trajectory $S^{(-\infty,\infty)}x$ is sufficient, which means, by definition, that some of its bounded segments $S^{[a,b]}x$ are sufficient.

In the case of an orbit $S^{(-\infty,\infty)}x$ with a simple singularity, sufficiency means that both branches of $S^{(-\infty,\infty)}x$ are sufficient.

The great importance of the sufficiency of a phase point x_0 is given by the fact that (for the discrete time billiard map, i. e. for the first return map T to $\partial\mathbf{M}$) such a phase point always has a suitably small, neighborhood $U_0 \ni x_0$ for which the first return map of T to U_0 exhibits a uniformly hyperbolic behavior, see Lemma 2.13 in [K-S-Sz(1990)].

For a more detailed exposition of the above notions and facts, the reader is kindly referred to §2 of [K-S-Sz(1990)].

No accumulation (of collisions) in finite time. By the results of Vaserstein [V(1979)], Galperin [G(1981)] and Burago-Ferleger-Kononenko [B-F-K(1998)], in a semi-dispersive billiard flow there can only be finitely many collisions in finite time intervals, see Theorem 1 in [B-F-K(1998)]. Thus, the dynamics is well defined as long as the trajectory does not hit more than one boundary components at the same time.

§3. THE IMPROVEMENTS

Let us see – step by step – the additions to the proof of Theorem 3.6 (of [K-S-Sz(1990)]) that facilitate the dropping of the requirement for the algebraic nature of the smooth components $\partial\mathbf{Q}_i$ from the set of conditions for the Theorem on Local Ergodicity. Our main reference in this section will be the mentioned paper. We try to follow its notations as closely as we can.

As it was said in 4. of [K-S-Sz(1991)], throughout the whole proof of Theorem 3.6 (of [K-S-Sz(1990)]) one has to use the so called “tubular distance function” $z(x) = z_{\text{tub}}(x)$ to appropriately measure the distance between the phase point $x = (q, v) \in \mathbf{M} \setminus \partial\mathbf{M}$ and the set of singular reflections $\mathcal{SR}$. This function $z(x) = z_{\text{tub}}(x)$ is to take the role of the function $\rho(x)$ in the original proof. Let us briefly recall here the definition of $z(x)$ for any $x = (q, v) \in \mathbf{M} \setminus \partial\mathbf{M}$:

For any phase point $y \in \mathbf{M} \setminus \partial\mathbf{M}$ introduce the number

$$t^+(y) = \min \{t > 0 \mid S^t y \in \partial\mathbf{M}\}.$$

The function $t^-(y)$ is defined as $-t^+(-y)$, where $q(-y) = q(y)$, and $v(-y) = -v(y)$.

Recall that the functions $t^+(\cdot)$ and $t^-(\cdot)$ turn out to be uniformly bounded, thanks to the introduction of the so called “virtual walls” (or “transparent walls”) to the model by Chernov and Sinai in [S-Ch(1987)], see also the second paragraph on p. 539 of [K-S-Sz(1990)]. These virtual walls make the system having a finite horizon. Caution: In order to keep the validity of condition (2) of the theorem, we have to drop (from the billiard flow) all trajectories gliding along a virtual wall over some time interval. The set of these dropped phase points has codimension two (a “slim set”), so it is negligible in all dynamical considerations.

The function $z(x)$ is then defined as

$$z(x) = \sup \left\{ r > 0 \mid S^{t^+(y)}y \notin \mathcal{SR} \text{ and } S^{t^-(y)}y \notin \mathcal{SR} \right. \\ \left. \text{for all } y \in \mathbf{M} \text{ with } v(y) = v(x), q(y) - q(x) \perp v(x), \|q(y) - q(x)\| < r \right\}.$$

(Here the supremum of the empty set is considered to be zero.)

We note that throughout this paper we use the Riemannian phase distance $d(x, y)^2 = \|q(x) - q(y)\|^2 + \|v(x) - v(y)\|^2$ to measure the distances between phase points $x, y \in \mathbf{M}$.

The invertible, measure-preserving mapping $T : \partial\mathbf{M} \rightarrow \partial\mathbf{M}$ that we study is now the first return map from $\partial\mathbf{M}$ to $\partial\mathbf{M}$. Its natural invariant measure μ_1 can be obtained from the Liouville measure μ of the flow $\{S^t\}_{t \in \mathbb{R}}$ by projection to $\partial\mathbf{M}$, i. e.

$$\mu_1(A) = \lim_{\epsilon \searrow 0} \epsilon^{-1} \mu(\{S^t y \mid y \in A, 0 \leq t \leq \epsilon\}),$$

for any Borel measurable subset A of $\partial\mathbf{M}$.

Lemma 1. For every fixed exponent $n \in \mathbb{N}$ the open δ -neighborhood

$$B_\delta(\mathcal{R}^{\leq n}) = \{x \in \partial\mathbf{M} \mid d(x, \mathcal{R}^{\leq n}) < \delta\}$$

of the compact singularity set

$$\mathcal{R}^{\leq n} = \bigcup_{k=1}^n \mathcal{R}^k = \{y \in \partial\mathbf{M} \mid T^n \text{ is not smooth on any neighborhood of } y \text{ in } \partial\mathbf{M}\}$$

of T^n has μ_1 -measure $O(\delta)$ (big ordo of δ), i. e. $\mu_1(B_\delta(\mathcal{R}^{\leq n})) \leq c_1 \delta$ with a positive constant c_1 depending only on the billiard flow, and n . (By the way, this convention applies to all constants c_i in this paper! The overbar denotes the closure taken in $\partial\mathbf{M}$.)

Proof. We start the proof with an introduction to the time evolution of the normal vector of a smooth, (locally) flow invariant, codimension-one submanifold J .

Given a codimension-one, flow-invariant, smooth sub-manifold $J \subset \mathbf{M}$, consider a normal vector $n_0 = (z, w)$ ($\neq 0$) of J at the phase point $y \in J$, i. e. for any tangent vector $(\delta q, \delta v) \in \mathcal{T}_y \mathbf{M}$ the relation $(\delta q, \delta v) \in \mathcal{T}_y J$ is true if and only if $\langle \delta q, z \rangle + \langle \delta v, w \rangle = 0$. Here $\langle \cdot, \cdot \rangle$ is the Euclidean inner product of the tangent space $\mathbb{R}^d$ of $\mathbf{Q}$ at every point $q \in \mathbf{Q}$. Let us determine first the time-evolution $n_0 \mapsto n_t$ ($t > 0$) of this normal vector as time t elapses. If there is no collision on the orbit segment $S^{[0,t]}y$, then the relationship between $(\delta q, \delta v) \in \mathcal{T}_y \mathbf{M}$ and $(\delta q', \delta v') = (DS^t)(\delta q, \delta v)$ is obviously

$$(3.1) \quad \begin{aligned} \delta v' &= \delta v, \\ \delta q' &= \delta q + t\delta v, \end{aligned}$$

from which we obtain that

$$\begin{aligned} (\delta q', \delta v') \in \mathcal{T}_{y'} J &\Leftrightarrow \langle \delta q' - t\delta v', z \rangle + \langle \delta v', w \rangle = 0 \\ &\Leftrightarrow \langle \delta q', z \rangle + \langle \delta v', w - tz \rangle = 0. \end{aligned}$$

This means that $n_t = (z, w - tz)$. It is always very useful to consider the quadratic form $Q(n) = Q((z, w)) =: \langle z, w \rangle$ associated with the normal vector $n = (z, w) \in \mathcal{T}_y \mathbf{M}$ of J at y . $Q(n)$ is the so called “infinitesimal Lyapunov function”, see [K-B(1994)] or part A.4 of the Appendix in [Ch(1994)]. For a detailed exposition of the relationship between the quadratic form Q , the relevant symplectic geometry and the dynamics, please see [L-W(1995)].

Remark. Since the normal vector $n = (z, w)$ of J is only determined up to a nonzero scalar multiplier, the value $Q(n)$ is only determined up to a positive multiplier. However, this means that the sign of $Q(n)$ (which is the utmost important thing for us) is uniquely determined. This remark will gain a particular importance in the near future.

From the above calculations we get that

$$(3.2) \quad Q(n_t) = Q(n_0) - t||z||^2 \leq Q(n_0), \quad (t > 0).$$

The next question is how the normal vector n of J gets transformed $n^- \mapsto n^+$ through a collision (reflection) at time $t = 0$? Elementary geometric considerations show (see Lemma 2 of [Sin(1979)], or formula (2) in §3 of [S-Ch(1987)]) that the linearization of the flow

$$(DS^t) \Big|_{t=0} : (\delta q^-, \delta v^-) \mapsto (\delta q^+, \delta v^+)$$

is given by the formulas

$$(3.3) \quad \begin{aligned} \delta q^+ &= R\delta q^-, \\ \delta v^+ &= R\delta v^- + 2\cos\phi RV^*KV\delta q^-, \end{aligned}$$

where the operator $R : \mathcal{T}_q\mathbf{Q} \rightarrow \mathcal{T}_q\mathbf{Q}$ is the orthogonal reflection across the tangent hyperplane $\mathcal{T}_q\partial\mathbf{Q}$ of $\partial\mathbf{Q}$ at $q \in \partial\mathbf{Q}$ ($y^- = (q, v^-) \in \partial\mathbf{M}$, $y^+ = (q, v^+) \in \partial\mathbf{M}$), $V : (v^-)^\perp \rightarrow \mathcal{T}_q\partial\mathbf{Q}$ is the v^- -parallel projection of the orthocomplement hyperplane $(v^-)^\perp$ onto $\mathcal{T}_q\partial\mathbf{Q}$, $V^* : \mathcal{T}_q\partial\mathbf{Q} \rightarrow (v^-)^\perp$ is the adjoint of V , i. e. it is the projection of $\mathcal{T}_q\partial\mathbf{Q}$ onto $(v^-)^\perp$ being parallel to the normal vector $\nu(q)$ of $\partial\mathbf{Q}$ at $q \in \partial\mathbf{Q}$, $K : \mathcal{T}_q\partial\mathbf{Q} \rightarrow \mathcal{T}_q\partial\mathbf{Q}$ is the second fundamental form of $\partial\mathbf{Q}$ at q and, finally, $\cos\phi = \langle \nu(q), v^+ \rangle$ is the cosine of the angle ϕ subtended by v^+ and the normal vector $\nu(q)$. For the formula (3.3), please also see the last displayed formula of §1 in [S-Ch(1982)], or (i) and (ii) of Proposition 2.3 in [K-S-Sz(1990)]. We note that it is enough to deal with the tangent vectors $(\delta q^-, \delta v^-) \in (v^-)^\perp \times (v^-)^\perp$ ($(\delta q^+, \delta v^+) \in (v^+)^\perp \times (v^+)^\perp$), for the manifold J under investigation is supposed to be flow-invariant, so any vector $(\delta q, \delta v) = (\alpha v, 0)$ ($\alpha \in \mathbb{R}$) is automatically inside $\mathcal{T}_y J$. The backward version (inverse)

$$(DS^t) \Big|_{t=0} : (\delta q^+, \delta v^+) \mapsto (\delta q^-, \delta v^-)$$

can be deduced easily from (3.3):

$$(3.4) \quad \begin{aligned} \delta q^- &= R\delta q^+, \\ \delta v^- &= R\delta v^+ - 2\cos\phi RV_1^*KV_1\delta q^+, \end{aligned}$$

where $V_1 : (v^+)^\perp \rightarrow \mathcal{T}_q\partial\mathbf{Q}$ is the v^+ -parallel projection of $(v^+)^\perp$ onto $\mathcal{T}_q\partial\mathbf{Q}$. By using formula (3.4), one easily computes the time-evolution $n^- \mapsto n^+$ of a normal vector $n^- = (z, w) \in \mathcal{T}_{y^-}\mathbf{M}$ of J if a collision $y^- \mapsto y^+$ takes place at time $t = 0$:

$$\begin{aligned} (\delta q^+, \delta v^+) \in \mathcal{T}_{y^+}J &\Leftrightarrow \langle R\delta q^+, z \rangle + \langle R\delta v^+ - 2\cos\phi RV_1^*KV_1\delta q^+, w \rangle = 0 \\ &\Leftrightarrow \langle \delta q^+, Rz - 2\cos\phi V_1^*KV_1Rw \rangle + \langle \delta v^+, Rw \rangle = 0. \end{aligned}$$

This means that

$$(3.5) \quad n^+ = (Rz - 2\cos\phi V_1^*KV_1Rw, Rw)$$

if $n^- = (z, w)$. It follows that

$$\begin{aligned}
(3.6) \quad Q(n^+) &= Q(n^-) - 2 \cos \phi \langle V_1^* K V_1 R w, R w \rangle \\
&= Q(n^-) - 2 \cos \phi \langle K V_1 R w, V_1 R w \rangle \leq Q(n^-).
\end{aligned}$$

Here we used the fact that the second fundamental form K of $\partial \mathbf{Q}$ at q is positive semi-definite, which just means that the billiard system is semi-dispersive.

The last, simple observation on the quadratic form $Q(n)$ regards the involution $I : \mathbf{M} \rightarrow \mathbf{M}$, $I(q, v) = (q, -v)$ corresponding to the time reversal. If $n = (z, w)$ is a normal vector of J at y then, obviously, $I(n) = (z, -w)$ is a normal vector of $I(J)$ at $I(y)$ and

$$(3.7) \quad Q(I(n)) = -Q(n).$$

Singularities.

Consider a smooth, connected piece $\mathcal{S} \subset \mathbf{M}$ of a singularity manifold corresponding to a singular (tangential or double) reflection some time *in the past*. Such a manifold $\mathcal{S}$ is locally flow-invariant and has one codimension, so we can speak about its normal vectors n and the uniquely determined sign of $Q(n)$ for $0 \neq n \in \mathcal{T}_y \mathbf{M}$, $y \in \mathcal{S}$, $n \perp \mathcal{S}$ (depending on the foot point, of course). Consider first a phase point $y^+ \in \partial \mathbf{M}$ right after the singular reflection that is described by $\mathcal{S}$. It follows from the proof of Lemma 4.1 of [K-S-Sz(1990)] and Sub-lemma 4.4 therein that at $y^+ = (q, v^+) \in \partial \mathbf{M}$ any tangent vector $(0, \delta v) \in \mathcal{T}_{y^+} \mathbf{M}$ lies actually in $\mathcal{T}_{y^+} \mathcal{S}$ and, consequently, the normal vector $n = (z, w) \in \mathcal{T}_{y^+} \mathbf{M}$ of $\mathcal{S}$ at y^+ necessarily has the form $n = (z, 0)$, especially $w = 0$. Thus $Q(n) = 0$ for any normal vector $n \in \mathcal{T}_{y^+} \mathbf{M}$ of $\mathcal{S}$. According to the monotonicity inequalities (3.2) and (3.6) above,

$$(3.8) \quad Q(n) < 0$$

for any phase point $y \in \mathcal{S}$ of a past singularity manifold $\mathcal{S}$.

For every $x \in \partial \mathbf{M}$ let $\Sigma_x \subset \partial \mathbf{M}$ be the following convex, local orthogonal manifold (projected on $\partial \mathbf{M}$ by the flow) containing x :

$$\begin{aligned}
(3.9) \quad \Sigma_x &= \{y \in \partial \mathbf{M}_i \mid \exists a, b > 0 \text{ such that } S^{(0,a]} x \cap \partial \mathbf{M} = \emptyset = S^{(0,b]} y \cap \partial \mathbf{M}, \\
&\quad q(S^b y) - q(S^a x) = (1+a)(v^+(y) - v^+(x))\},
\end{aligned}$$

where $\partial \mathbf{M}_i$ is the smooth component of the boundary $\partial \mathbf{M}$ containing x . It is clear that for every given $a > 0$ (with $S^{(0,a]} x \cap \partial \mathbf{M} = \emptyset$) the “synchronized wave front”

$$\begin{aligned} \Sigma_x^a &= \{S^b y \in \partial \mathbf{M}_i \mid b > 0, S^{(0,b]} y \cap \partial \mathbf{M} = \emptyset, \\ & q(S^b y) - q(S^a x) = (1+a) (v^+(y) - v^+(x)) \} \end{aligned}$$

is a sphere (in the configuration space $\mathbf{Q}$) of radius $1+a$ supplied with the field of unit outer normal vectors containing the point $q(S^a x)$. Its backward projection (along the flow) onto $\partial \mathbf{M}$ is Σ_x . In the simple calculations below, leading to the proof of Proposition 3.10, all objects are considered *after the collision* taking place at a phase point in $\partial \mathbf{M}$, especially Σ_x turns out to be a spherical, local, convex orthogonal manifold. The curvature operator (second fundamental form) of $S^a(\Sigma_x)$ at the point $S^b y$ (here a and b are the same numbers as in (3.9)) is $(1+a)^{-1} \cdot I$, which operator is uniformly positive. (We can restrict ourselves to the case $0 < a \leq 1$.) On the other hand, as (3.7) and (3.8) show, the infinitesimal Lyapunov function $Q(n) = \langle z, w \rangle$ is positive for the flow-invariant hull J of any future singularity. Let $y' \in J$, $n = (z, w)$ a unit normal vector of J at y' , and for any $y' = S^b y \in S^a(\Sigma_x)$ ($x \in \partial \mathbf{M}$, a and b are the same numbers as in (3.9)) define the unit tangent vector $\vec{\tau} = (\delta q, (1+a)^{-1} \delta q) \in \mathcal{T}_{y'} S^a(\Sigma_x)$ in such a way that the vector δq has the same direction as the vector $z + (1+a)^{-1} w$, i. e.

$$\delta q = (1 + (1+a)^{-2})^{-1/2} \|z + (1+a)^{-1} w\|^{-1} (z + (1+a)^{-1} w).$$

Then for the scalar product of the unit vectors $\vec{\tau}$ and n we get the following inequalities:

$$\begin{aligned} \langle \delta q, z \rangle + (1+a)^{-1} \langle \delta q, w \rangle &= \|\delta q\| \cdot \|z + (1+a)^{-1} w\| \\ &\geq (1 + (1+a)^{-2})^{-1/2} \cdot \min \{1, (1+a)^{-1}\} \geq \text{const} > 0 \end{aligned}$$

The uniform positivity of the scalar product of the tangent vector $\vec{\tau}$ of Σ_x and the normal vector n of J proves our

Proposition 3.10. The manifolds Σ_x are uniformly transversal to the singularity components $K_i \subset \mathcal{R}^{\leq n}$ of T^n .

Continuing the proof of Lemma 1. Thanks to the compactness of the phase space $\partial \mathbf{M}$, it is enough to prove that any given phase point $x_0 \in \partial \mathbf{M}$ has a neighborhood U_0 in $\partial \mathbf{M}$ such that the estimation of Lemma 1 holds true for the singularity set $U_0 \cap \mathcal{R}^{\leq n}$. In the proof of Lemma 1 we will always be dealing with the post-collision velocities v^+ of phase points $x = (q, v^+) \in \partial \mathbf{M}$. In order to slightly simplify the notations, we assume that $x_0 \notin \partial \partial \mathbf{M}$. The proof in the case $x_0 \in \partial \partial \mathbf{M}$ can be obtained from the one below by small and straightforward modifications.

According to Proposition 3.10 and its proof, the phase point $x_0 \in \text{int} \partial \mathbf{M}$ has a closed neighborhood $\tilde{U}_0$ in $\partial \mathbf{M}$ and a smooth coordinate patch $\Phi : B_{\epsilon_0} \times [-\epsilon_1, \epsilon_1] \xrightarrow{\cong} \tilde{U}_0$ (here $B_{\epsilon_0} = \{y \in \mathbb{R}^{2d-3} \mid \|y\| \leq \epsilon_0\}$) with the following properties:

(1) $\Phi(0, 0) = x_0$;

(2) For the parametrized curves γ_y , $\gamma_y(t) = \Phi(y, t)$ ($y \in B_{\epsilon_0}$), the following is true: If $\gamma_y(t) \in \text{int} K_i$, α is the non-obtuse angle between the tangent vector $\dot{\gamma}_y(t)$ and the normal vector $n_{\gamma_y(t)}(K_i)$ of K_i at $\gamma_y(t)$, then $\cos \alpha \geq c_2$ (> 0) with a constant c_2 depending only on the model and n ;

(3) $\sup_{y \in B_{\epsilon_0}} |\gamma_y \cap \mathcal{R}^{\leq n}| < \infty$, where $|\cdot|$ denotes the cardinality of a set.

The construction of the neighborhood $\tilde{U}_0$ of x_0 (fulfilling (1)–(3)) goes on as follows: First we select the curve $\gamma_0 \ni x_0$ as the intersection $\Sigma_{x_0} \cap W_{x_0}$ in such a way that γ_0 be transversal to all components K_i of $\mathcal{R}^{\leq n}$ near x_0 . Here

$$W_y = \{(q, v^+) \in \partial \mathbf{M} \mid v^+ = \lambda v^+(y) + \mu w_0, \quad \lambda, \mu \in \mathbb{R}\},$$

for any phase point $y \in \partial \mathbf{M}$ near x_0 , where $w_0 \in \mathbb{R}^d$ is a suitably selected (and fixed) vector which is not parallel to the velocity $v^+(x_0)$. Then we construct the smooth, codimension-one manifold $B_0 = \Phi(B_{\epsilon_0} \times \{0\}) \ni x_0$ near x_0 in such a way that B_0 is perpendicular to γ_0 at x_0 . Finally, the other fibers γ_y ($y \in B_{\epsilon_0}$) are constructed as $\gamma_y = \Sigma_{y'} \cap W_{y'}$, where $y' = \Phi(y, 0) \in B_0$, and $W_{y'}$ is defined above. By applying Proposition 3.10, we can make it sure that the parametrized set $\tilde{U}_0 = \Phi(B_{\epsilon_0} \times [-\epsilon_1, \epsilon_1])$ (with a suitably selected, small $\epsilon_1 > 0$) is, indeed, a closed neighborhood of x_0 with the properties (1)–(2). The validity of the uniform finiteness (3) follows from the fact that the billiard flow has a finite horizon (with the introduced virtual walls), thus the n -th iterate T^n of the billiard map $T : \partial \mathbf{M} \rightarrow \partial \mathbf{M}$ can only break the local, convex curves $\gamma_y = \Sigma_{y'} \cap W_{y'}$ into finitely many connected components, where “finite” is meant to be uniformly finite in y for a fixed n .

To finish the proof of Lemma 1, we need our

Sub-lemma. Given the above triplet $(x_0, \tilde{U}_0, \Phi)$, we can find a number $\epsilon_2 > 0$, $\epsilon_2 < \epsilon_1$, such that for any small enough $\delta > 0$ ($\delta < \delta_0$) and for any phase point $z \in U_0 := \Phi(B_{\epsilon_0} \times [-\epsilon_2, \epsilon_2])$, $z = \Phi(y, t)$ ($y \in B_{\epsilon_0}$, $|t| \leq \epsilon_2$), the inequality $d(z, \mathcal{R}^{\leq n}) < \delta$ implies that there exists a number $t' \in [-\epsilon_1, \epsilon_1]$, $|t - t'| \leq 2c_2^{-1}\delta$, $\Phi(y, t') \in \mathcal{R}^{\leq n} \cap \tilde{U}_0$.

Proof. We may assume that the parametrization t along γ_y (i. e. $\gamma_y(t) = \Phi(y, t)$) is asymptotically the same as the Riemannian distance in $\tilde{U}_0$. The statement of the sub-lemma is a direct consequence of the transversality property (2) and the well known fact that any smooth component K_i of $\mathcal{R}^{\leq n}$ can only terminate on another component K_j of $\mathcal{R}^{\leq n}$ having actually smaller degree than K_i . (The degree k of a component K_i of $\mathcal{R}^{\leq n}$ is the uniquely determined positive integer k for which $K_i \subset T^{-k}(\mathcal{SR}^-)$.) For varying $z \in \gamma_y$, $z = \Phi(y, t)$, we take the level set $L_t = \{\Phi(y', t) \mid y' \in B_{\epsilon_0}\}$, and consider a closest point $r(z)$ of $L_t \cap \mathcal{R}^{\leq n}$ to z . It follows from (2) that in one direction on γ_y the distance function $d(z, r(z))$ decreases

at least at the rate of c_2^{-1} with respect to the parameter t . In the requirement $|t - t'| \leq 2c_2^{-1}\delta$ we included the additional factor 2 just to be on the safe side, for the parametrization of γ_y is only asymptotically the same as the Riemannian distance in $\tilde{U}_0$. $\square$

We may assume that in the neighborhood $\tilde{U}_0$ the Radon-Nikodym derivative of μ_1 with respect to the Lebesgue measure λ (defined by the coordinate patch Φ) is between two constants close to 1 so, in any case, we have that $\mu_1(A) \leq c_3\lambda(A)$ for any Borel set $A \subset \tilde{U}_0$. Set $N_0 := \sup_{y \in B_{\epsilon_0}} |\gamma_y \cap \mathcal{R}^{\leq n}|$. According to the properties (1)–(3), the sub-lemma, and the above observation we have the chain of inequalities

$$\begin{aligned}
 (3.11) \quad & \mu_1(U_0 \cap B_\delta(\mathcal{R}^{\leq n})) \leq c_3\lambda\{z = \Phi(y, t) \mid y \in B_{\epsilon_0}, \\
 & |t| \leq \epsilon_2, \exists t' \in [-\epsilon_1, \epsilon_1], |t - t'| \leq 2c_2^{-1}\delta, \Phi(y, t') \in \mathcal{R}^{\leq n}\} \\
 & \leq c_3 \int_{B_{\epsilon_0}} 4c_2^{-1}\delta N_0 d\lambda(B_{\epsilon_0}) = 4c_2^{-1}\delta N_0 c_3 \text{Vol}(B_{\epsilon_0}) = O(\delta),
 \end{aligned}$$

and the proof of Lemma 1 is now complete. $\square$

Corollary 1. For every fixed exponent $n \in \mathbb{N}$ the set

$$A_n^\delta = \bigcup \{G_i^\delta \mid G_i^\delta \in \mathcal{G}^\delta, G_i^\delta \cap \mathcal{R}^{\leq n} \neq \emptyset\}$$

has μ_1 -measure big ordo of δ , i. e. $\mu_1(A_n^\delta) \leq c_1\delta$ with a positive constant c_1 depending only on the billiard flow and n . (For the definition of a sequence of regular coverings $\mathcal{G}^\delta = \{G_i^\delta \mid i \in I(\delta)\}$, $0 < \delta < \delta_0$, please see §3 of [K-S-Sz(1990)].)

We need to save the crucial geometric inequality of Lemma 4.6 of [K-S-Sz(1990)].

Lemma 2. For any fixed exponent $n \in \mathbb{N}$ denote by K_i ($i = 1, 2, \dots, N$) all the smooth components of the compact singularity set $K = \mathcal{R}^{\leq n}$ of T^n . We claim that the set

$$B_n^\delta = \{x \in \partial \mathbf{M} \mid \exists i < j \text{ such that } d(x, K_i) < \delta, d(x, K_j) < \delta\}$$

has μ_1 -measure $o(\delta)$ (small ordo of δ).

Proof. The proof is a beefed-up version of the proof of the previous lemma.

By repeating the proof of Lemma 1, reaching its climax in (3.11), we see that for every phase point $z = \Phi(y, t) \in U_0 \cap B_n^\delta$ ($y \in B_{\epsilon_0}$, $|t| \leq \epsilon_2$) one can find a $t' \in [-\epsilon_1, \epsilon_1]$ for which $|t - t'| \leq 2c_2^{-1}\delta$, $z' = \Phi(y, t') \in K_i \cap \tilde{U}_0$, $d(z', K_j) \leq c_4\delta$, and K_i, K_j are two distinct smooth components of $\mathcal{R}^{\leq n}$. Observe that for any point $z' \in K_i \setminus \bigcup_{j=1}^k \partial K_j$ the distance between z' and any K_j (with $j \neq i$) is positive. (Here $\mathcal{R}^{\leq n} = \bigcup_{j=1}^k K_j$ is the canonical subdivision of the singularity set $\mathcal{R}^{\leq n}$ into

its smooth components.) We conclude that for the phase point $z' = \Phi(y, t')$ above (constructed for the point $z = \Phi(y, t) \in U_0 \cap B_n^\delta$) a distance estimation

$$d\left(z', \bigcup_{j=1}^k \partial K_j\right) \leq f(\delta)$$

holds true with a function $f(\delta) > 0$, for which $\lim_{\delta \rightarrow 0} f(\delta) = 0$. Set

$$A_\delta = \left\{ y \in B_{\epsilon_0} \mid \exists t' \in [-\epsilon_1, \epsilon_1] \text{ such that } d\left(\Phi(y, t'), \bigcup_{j=1}^k \partial K_j\right) \leq f(\delta) \right\}.$$

Since the intersection $\bigcap_{\delta > 0} A_\delta$ is the obviously zero-measured set

$$\left\{ y \in B_{\epsilon_0} \mid \exists t' \in [-\epsilon_1, \epsilon_1] \text{ such that } \Phi(y, t') \in \bigcup_{j=1}^k \partial K_j \right\},$$

we get that $\lim_{\delta \rightarrow 0} \text{Vol}(A_\delta) = 0$, where $\text{Vol}(\cdot)$ is the $(2d-3)$ -dimensional Lebesgue measure in B_{ϵ_0} . Therefore, the estimation (3.11) now gives us the upper bound

$$\mu_1(U_0 \cap B_n^\delta) \leq c_3 \int_{A_\delta} 4c_2^{-1} \delta N_0 d\lambda(B_{\epsilon_0}) = 4c_2^{-1} \delta N_0 c_3 \text{Vol}(A_\delta) = o(\delta),$$

which finishes the proof of Lemma 2. $\square$

Corollary 2. The set

$$\{x \in U_0 \mid \exists i < j \exists G_k^\delta \in \mathcal{G}^\delta \text{ such that } x \in G_k^\delta, G_k^\delta \cap K_i \neq \emptyset \neq G_k^\delta \cap K_j\}$$

has μ_1 -measure $o(\delta)$ (small ordo of δ).

This has been the last step in the revision of the proof of the Theorem on Local Ergodicity for semi-dispersive billiards.

Acknowledgment. The author is deeply indebted to N. Chernov for his making a great deal of very useful, enlightening remarks.

REFERENCES

- [B-Ch-Sz-T(2002)] P. Bálint, N. Chernov, D. Szász, I. P. Tóth, *Multidimensional semidispersing billiards: singularities and the fundamental theorem*, Ann. Henri Poincaré **3** no. **3**, 451–482.
- [B-F-K(1998)] D. Burago, S. Ferleger, A. Kononenko, *A geometric approach to semi-dispersing billiards*, Ergod. Th. & Dynam. Sys. **18**, 303–319.
- [Ch(1994)] N. I. Chernov, *Statistical Properties of the Periodic Lorentz Gas. Multidimensional Case*, Journal of Statistical Physics **74**, Nos. **1/2**, 11–54.
- [G(1981)] G. Galperin, *On systems of locally interacting and repelling particles moving in space*, Trudy MMO **43**, 142–196.
- [K-B(1994)] A. Katok, K. Burns, *Infinitesimal Lyapunov functions, invariant cone families and stochastic properties of smooth dynamical systems*, Ergodic Theory Dyn. Syst. **14**, No. **4**, 757–785.
- [K-S-Sz(1990)] A. Krámli, N. Simányi, D. Szász, *A “Transversal” Fundamental Theorem for Semi-Dispersing Billiards*, Commun. Math. Phys. **129**, 535–560.
- [K-S-Sz(1991)] A. Krámli, N. Simányi, D. Szász, *Erratum. A “Transversal” Fundamental Theorem for Semi-Dispersing Billiards*, Commun. Math. Phys. **138**, 207–208.
- [L-W(1995)] C. Liverani, M. Wojtkowski, *Ergodicity in Hamiltonian systems*, Dynamics Reported **4**, 130–202, arXiv:math.DS/9210229..
- [Sim(2002)] N. Simányi, *The Complete Hyperbolicity of Cylindric Billiards*, Ergod. Th. & Dynam. Sys. **22**, 281–302, arXiv:math.DS/9906139.
- [Sin(1979)] Ya. G. Sinai, *Development of Krylov’s ideas. Afterword to N. S. Krylov’s “Works on the foundations of statistical physics”, see reference [K(1979)]*, Princeton University Press.
- [S-Ch(1982)] Ya. G. Sinai, N.I. Chernov, *Entropy of a gas of hard spheres with respect to the group of space-time shifts*, Trudy Sem. Petrovsk. No. **8**, 218–238.
- [S-Ch(1987)] Ya. G. Sinai, N.I. Chernov, *Ergodic properties of certain systems of 2-D discs and 3-D balls*, Russian Math. Surveys (**3**) **42**, 181–207.
- [S-Sz(1999)] N. Simányi, D. Szász, *Hard Ball Systems Are Completely Hyperbolic*, Annals of Math. **149**, 35–96, arXiv:math.DS/9704229..
- [S-Sz(2000)] N. Simányi, D. Szász, *Non-integrability of Cylindric Billiards and Transitive Lie Group Actions*, Ergod. Th. & Dynam. Sys. **20**, 593–610.
- [V(1979)] L. N. Vaserstein, *On Systems of Particles with Finite Range and/or Repulsive Interactions*, Commun. Math. Phys. **69**, 31–56.